

General information about positive NIPT results: Trisomy 10

My patient's NIPT is positive for trisomy 10. What does this mean?

Your patient's noninvasive prenatal testing (NIPT) result suggests the presence of an extra copy of chromosome 10. NIPT is a screening test; false positives can occur. The actual chance for the pregnancy to have trisomy 10 depends on many factors, including the patient's clinical and family history.

Next steps to consider: You should discuss the results and the potential clinical implications with your patient. Globally, professional medical societies recommend that all women with a positive screening result should have genetic counseling and a comprehensive ultrasound evaluation with an opportunity for diagnostic testing to confirm the results.^{1,2} Confirmation prior to birth can also help with pregnancy and neonatal management.

See below for more information about trisomy 10.

What is trisomy 10?

Trisomy 10 is a condition that is caused by an extra chromosome number 10 (three copies instead of two).

What are the features of trisomy 10?

Most pregnancies with trisomy 10 will miscarry spontaneously. If a developing fetus has mosaic trisomy 10 (where some cells are normal and some cells have trisomy 10), there is an increased chance for the pregnancy to progress and possibly survive to term. However, liveborn infants with full or mosaic trisomy 10 are expected to have serious medical problems. Key features include dysmorphism, cardiac and renal anomalies, growth retardation, and deformities of hands and feet. Neonatal or early infancy death is the most likely outcome. In reported cases of prenatally diagnosed trisomy 10, the outcomes have ranged from normal to live births with clinical sequelae. The variability in clinical presentation is believed to be due to confined placental mosaicism (CPM; when trisomic cells are present in the placenta, but not in the fetus) or the degree of fetal mosaicism.

What is the prevalence of this condition?

Unknown, but rare. For this reason, positive predictive value (PPV) cannot be accurately calculated.

What testing could be considered?

Specialized genetic tests such as karyotyping, fluorescence *in situ* hybridization (FISH), quantitative polymerase chain reaction (qPCR), and microarray are available to confirm the presence of trisomy 10.

These confirmatory tests are generally performed on cells from chorionic villus sampling (CVS) or amniocentesis during pregnancy, on cord blood or peripheral blood sample after the baby is born, or on products of conception (POC) in the case of a miscarriage. The type of invasive procedure and diagnostic testing should take into account the underlying chromosome anomaly.^{3,4}

Ultrasound evaluation may be useful in aiding with a prenatal diagnosis of trisomy 10, but a normal ultrasound cannot exclude trisomy 10.

Resources for trisomy 10

MedlinePlus Genetics

medlineplus.gov/genetics/chromosome/10

Unique, The Rare Chromosome Disorder Support Group

rarechromo.org

References

- Rose NC, Kaimal AJ, Dugoff L, et al. Screening for Fetal Chromosomal Abnormalities: ACOG Practice Bulletin, Number 226. *Obstet Gynecol.* 2020;136(4):e48-e69.
- Benn P, Borrell A, Chiu RW, et al. Position statement from the Chromosome Abnormality Screening Committee on behalf of the Board of the International Society for Prenatal Diagnosis. *Prenat Diagn.* 2015;35(8):725-734.
- Cherry AM, Akkari YM, Barr KM, et al. Diagnostic cytogenetic testing following positive noninvasive prenatal screening results: a clinical laboratory practice resource of the American College of Medical Genetics and Genomics (ACMG). *Genet Med.* 2017;19(8):845-850.
- Van Opstal D, Srebniak MI. Cytogenetic confirmation of a positive NIPT result: evidence-based choice between chorionic villus sampling and amniocentesis depending on chromosome aberration. *Expert Rev Mol Diagn.* 2016;16(5):513-520.

Additional resources

Gardner RJM, Sutherland GR, Shaffer LG. *Gardner and Sutherland's Chromosome Abnormalities and Genetic Counseling.* 5th ed. Oxford University Press; 2018.

Gao Y, Ma YC, Ju YH, Li YN. Mosaicism trisomy 10 in a 14-month-old child with additional neurological abnormalities: case report and literature review. *BMC Pediatr.* 2018;18(1):266.

Lévy J, Jouannic JM, Saada J, Dhombres F, Siffroi JP, Portnoi MF. Prenatal diagnosis of bilateral ectrodactyly and radial agenesis associated with trisomy 10 mosaicism. *Case Rep Genet.* 2013;2013:592702.

Malvestiti F, Agrati C, Grimi B, et al. Interpreting mosaicism in chorionic villi: results of a monocentric series of 1001 mosaics in chorionic villi with follow-up amniocentesis. *Prenat Diagn.* 2015;35(11):1117-1127.